


Ena

**PALM BEACH COUNTY, FLORIDA  
DIVISION OF ENVIRONMENTAL  
SCIENCE AND ENGINEERING  
AIR POLLUTION CONTROL**

# **ANNUAL REPORT 1983**





## TABLE OF CONTENTS

ii	List of Tables and Figures
----	----------------------------

SECTION		PAGE
I.	Introduction.....	1
II.	Meteorology.....	6
III.	Technical Studies.....	13
IV.	Air Quality Index.....	52
V.	Emission Inventory.....	55



# LIST OF TABLES & FIGURES

		Page
Table 1	Population, Palm Beach County.....	5
Table 2	Meteorological Means.....	9
Table 3	Wind Direction & Speed - West Palm Beach.....	10
Table 4	Wind Direction & Speed - Royal Palm Beach.....	11
Table 5	Wind Direction & Speed - 20 Mile Bend.....	12
Table 6	Monitoring Site Locations.....	21
Figure 1	Ambient Air Monitoring Network.....	24
Table 7	Ambient Air Quality Standards.....	25
Table 8	Network Description.....	26
Table 9	Suspended Particulate Matter 1983.....	27
Table 10	Total Suspended Particulate, 1983.....	28
Figure 2	Range of Probable Logarithmic Values High Volume Samples, 1982-1983.....	29
Table 11	Total Suspended Particulate, 1969-1983.....	30
Table 12	Gaseous Sampling 1970-1983.....	34
Table 13	Nitrogen Dioxide Data, Site 1, 1983.....	40
Table 14	Ambient Air, Nitrogen Dioxide, Site 1, 1983..	41
Table 15	Carbon Monoxide, Data Site 1, 1983.....	42
Table 16	Ambient Air, Carbon Monoxide, Site 1, 1983..	43
Table 17	Ozone Data, Site 16, 1983.....	44
Table 18	Ambient Air - Ozone, Site 16, 1983.....	45
Table 19	Ozone Data, Site 21, 1983.....	46
Table 20	Ambient Air - Ozone, Site 21, 1983.....	47
Table 21	Ozone Design Values, Site 16, 1980-1983.....	48
Table 22	Ozone Design Values, Site 21, 1973-1983.....	49
Table 23	Sulfur Dioxide Data, Site 22, 1983.....	50
Table 24	Ambient Air, Sulfur Dioxide, Site 22, 1983..	51
Table 25	Air Quality Index.....	54
Table 26	Summary of Air Pollution Emissions in Palm Beach County, 1983 (T/YR) - Point Sources.....	59
Table 27	Summary of Air Pollution Emission in Palm Beach County, 1983 (T/YR) - Area Sources.....	60



# I

## INTRODUCTION

Palm Beach County is located along the southeast coast of Florida, and is part of what is generally called the "Florida Gold Coast". The "Gold Coast" consists of four counties: Palm Beach, Broward, Dade and Monroe. Palm Beach County is geographically separated into two regions: one region is a highly urbanized coastal strip, stretching approximately forty-five (45) miles from Tequesta on the north, to Boca Raton on the south; and the other is agricultural, located in the western portion of Palm Beach County (The Glades Area).

The population distribution within the County depicts this same unusual pattern. Virtually all of the population is located in two relatively small areas: the coastal strip adjacent to the Atlantic Ocean and along a narrow band adjacent to Lake Okeechobee. A population table is included in this section (Table 1).

The majority of the population, approximately 90%, is located on the coast from a distance of approximately 15 miles inland from the Atlantic Ocean. Developments through the years have resulted in several municipalities along the coastal strip.

The band adjacent to Lake Okeechobee contains virtually all of the remaining 10% of the County's population. Municipalities which are included in the Glades area are: Belle Glade, Pahokee and South Bay. The economy of this area is an agricultural one, based principally on sugar cane and winter vegetables.



Tourism and related fields continue to be the major economic factors in the urban area. Tourism has increased along with increased population. Other major industries in the area include building construction and related fields, agriculture, aircraft testing facility, electronics, cement and concrete, asphaltic concrete and the service industries.

The complexity of the problems of Air Pollution Control are related to the widespread growth of Palm Beach County. Advances in environmental protection activities, which are being carried out by this program, have been utilized in order to keep abreast of Air Pollution problems. These activities are characteristic of urban areas across the nation.

The review of applications for the state air permits is one of the many activities handled by the Air Pollution Section of the Palm Beach County Health Department. The Florida Department of Environmental Regulation requires both a permit to construct and a permit to operate any air pollution source. The review of permit applications places our local program in a position to prevent the improper construction of a pollution source and to assure that adequate pollution control equipment is utilized and maintained.

Other activities include: consultations with industries and engineers on impending permit action; enforcement action; maintenance of monitoring network; and required compliance schedule and increments of progress surveillance. Also, our local program investigates and initiates the necessary follow-up action regarding all citizen compliants. As part of the State Air



Implementation Plan, this agency is required to conduct source inspections of existing and new sources in Palm Beach County, in order to assure that all sources are in compliance with Air Pollution Regulations.

The air monitoring capabilities of our program continues to be the ultimate means of maintaining air quality standards in Palm Beach County. The Environmental Control Air Monitoring Laboratory located in West Palm Beach is capable of continuously measuring stationary and mobile source related pollutants, i.e., Carbon Monoxide (CO), Nitrogen Dioxide (NO<sub>2</sub>), and four meteorological parameters (wind speed, wind direction, temperature, and relative humidity). The data is collected through the use of recorders plus a centrally controlled data acquisition system. Our ozone monitor was removed from the air monitoring laboratory in 1979 and relocated in the Town of Royal Palm Beach in order to comply with the E.P.A. siting criteria. The site is also equipped with a centrally controlled data acquisition system. A second ozone monitor was placed at our 20 Mile Bend Site. Both of the ozone sites are National Air Monitoring Sites (NAMS) and are equipped with continuous meteorological sensing equipment. Our sulfur dioxide (SO<sub>2</sub>) site was also relocated in accordance with E.P.A. criteria and placed at Riviera Beach.

In addition to the above, our air quality monitoring network includes ten (10) high volume particulate sites. All analysis pertaining to air pollution is performed by our chemistry laboratory located in Delray Beach. Furthermore, the program continues to monitor for the frequency and intensity of



temperature inversions at the Division of Forestry's observation tower located in Loxahatchee, Florida.

All data collected by our air monitoring network is transmitted via telephone lines using a video display system to the Department of Environmental Regulations in Tallahassee. The data is placed in SAROAD (Storage & Retrieval of Aerometric Data) format for transmission.

Public relation activities during the past year by the Air Pollution Section of the Palm Beach County Health Department have consisted of continued steps to inform the general public of the programs and procedures established to maintain our good air quality. These activities include wide distribution of our Annual Report, extending invitations to groups of school and environmental clubs to visit our facilities, and presentations to school and civic groups throughout the County on the topic of "Air Pollution Control in Palm Beach County". In addition, a major aspect of our program's public relations activities is the dissemination, twice daily, of an "Air Quality Index" to the local news media. The index utilizes the daily results of all measured pollutants.

TABLE 1

## POPULATION FOR PALM BEACH COUNTY AND MUNICIPALITIES

<u>MUNICIPALITY</u>	U.S. Census Bureau Counts 1980	Univ. of Fla. Population Estimates 1982	Univ. of Fla. Population Estimates 1983
Atlantis	1,325	1,517	1,530
Belle Glade	16,535	16,928	17,057
Boca Raton	49,505	51,302	52,315
Boynton Beach	35,624	37,532	38,102
Briny Breezes	387	381	384
Cloud Lake	160	145	148
Delray Beach	34,325	38,530	39,470
Glen Ridge	235	235	233
Golf	110	114	116
Golfview	210	205	208
Greenacres City	8,843	16,614	18,357
Gulfstream	475	481	498
Haverhill	1,249	1,257	1,255
Highland Beach	2,030	2,354	2,466
Hypoluxo	537	773	838
Juno Beach	1,142	1,241	1,221
Jupiter	9,868	13,274	14,917
Jupiter Inlet Colony	378	400	400
Lake Clarke Shores	3,174	3,184	3,170
Lake Park	6,909	6,900	6,871
Lake Worth	27,048	27,066	27,189
Lantana	8,048	8,355	8,387
Manalapan	329	348	355
Mangonia Park	1,419	1,377	1,367
North Palm Beach	11,344	12,005	12,123
Ocean Ridge	1,355	1,438	1,470
Pahokee	6,346	6,363	6,473
Palm Beach	9,729	10,391	10,499
Palm Beach Gardens	14,407	17,673	18,108
Palm Beach Shores	1,232	1,243	1,248
Palm Springs	8,166	9,146	9,171
Riviera Beach	26,596	26,634	26,573
Royal Palm Beach	3,423	4,712	5,177
South Bay	3,886	3,702	3,675
South Palm Beach	1,304	1,340	1,372
Tequesta	3,685	3,828	3,810
West Palm Beach	62,530	64,090	63,903
Total Incorporated	363,904	393,070	400,396
UNINCORPORATED	209,221	244,862	251,799
TOTAL COUNTY	573,125	637,940	652,195



## II

### METEOROLOGY

Topography is of primary importance whenever the meteorological aspects of a given region are to be evaluated. Palm Beach County is a fairly level region. For the most part, Palm Beach County is between 10 and 20 feet above sea level. All urban development is located along the eastern 15 miles of the coastal strip. Most of the western portions of the County are covered by agricultural lands or everglades.

The agricultural lands are endowed with a rich peatlike "muck" soil. The Atlantic Ocean borders the eastern edge of the County and the Gulf stream flows northward approximately 3 miles off-shore. Seldom does a cold air mass reach this region without being modified due to marine influences and our southern location. Light freezes occur infrequently along the coastal areas of the County and more frequently in the everglades and agricultural areas. The most eastern parts of the County come under the influence of the sea breeze during the day and land breeze during the night. Based on weather data accumulated at Palm Beach International Airport (Table 2), August is the warmest month with a mean of 82.7°F, a maximum mean of 90.6°F, and minimum mean of 74.7°F. From the same data, January is the coldest month with a mean of 65.9°F, a maximum mean of 75.1°F and a minimum mean of 56.7°F. Rain showers and/or thunderstorms of short duration are frequent during the summer season. Palm Beach County receives the greatest amount of rainfall during the

summer and fall. As indicated in Table 2, the County receives an average 60.9 inches of rain per year.

Palm Beach County can be classified as a semitropical region. The Quasi - permanent location of the "Bermuda" high pressure area governs our weather. It causes our prevailing easterly surface winds in addition to supplying the warm moist air necessary to produce the frequent air mass, frontal or nocturnal rain-showers and/or thunderstorms that occur in the County.

The position of the "Bermuda" high pressure area is also conducive to the formation of an atmosphere capable of causing high pollution days. This atmosphere can easily occur if cold air from the north moves underneath the warm moist air brought into the County by the "Bermuda" high. The result is a temperature inversion or increase of temperature with height which traps the pollutants in the lower levels.

Meteorological parameters play a significant role in understanding the over-all air pollution cycle. The motions of the atmosphere are extremely variable and must be thoroughly examined in order to determine the movement and dispersement of pollutants. Both wind direction and wind speed are of primary importance. The surface wind and the wind found in the first few hundred feet of our atmosphere must be studied to determine diffusion and movement of the pollutants. The wind direction is indicative of the direction of travel of the pollutants. The wind speed determines the time it takes the pollutants to travel to a receptor and is a function of the amount of dilution of pollutant. Light winds, coupled with other factors, contribute to poor air quality



episodes. U. S. Weather Service records of Palm Beach International Airport show the prevailing wind directions for the months of February through November are from one of the easterly headings. Mean monthly speeds vary between 7.6 mi/hr in August and 10.9 mi/hr in April (Table 2). The wind direction and speed for 1983 (Table 3) taken at our continuous monitoring site in West Palm Beach shows that a higher percentage of winds were from the east-northeast, east, or east southeast directions and the majority of the wind speeds were in the 4-10 mi/hr category. Table 4 & Table 5 wind direction and speed, for our Royal Palm Beach & 20 Mile Bend ozone sites (NAMS) are included in this section for information purposes.

TABLE 2

PALM BEACH INTERNATIONAL AIRPORT  
METEOROLOGICAL MEANS

MONTH	MEAN MAXIMUM TEMP.	MEAN MINIMUM TEMP.	MEAN MONTHLY TEMP.	MEAN MONTHLY PRECIP. (IN)	PREVAILING WIND DIRECTION	MEAN WIND SPEED (mi/hr)
JANUARY	75.1	56.7	65.9	2.64	NW	9.9
FEBRUARY	76.0	56.7	66.4	2.41	SE	10.3
MARCH	79.4	61.0	70.2	2.93	SE	10.8
APRIL	82.7	65.3	74.0	3.47	E	10.9
MAY	86.0	69.3	77.7	5.76	ESE	9.7
JUNE	88.7	72.6	80.7	7.89	ESE	8.1
JULY	90.4	74.3	82.4	6.37	ESE	7.5
AUGUST	90.6	74.7	82.7	6.52	ESE	7.6
SEPTEMBER	88.7	74.3	81.5	9.53	ENE	8.6
OCTOBER	84.8	70.4	77.6	7.74	ENE	10.0
NOVEMBER	79.9	63.9	71.9	3.14	ENE	10.1
DECEMBER	76.4	58.8	67.6	2.48	NNW	9.9
YEARLY	83.2	66.5	74.9	60.88	ESE	9.5



TABLE 3

SITE #1 - WEST PALM BEACH  
WIND DIRECTION AND SPEED (MI/HR) OCCURRENCES  
1983

DIRECTION	1-3	4-6	7-10	11-16	17-21	> 22	TOTAL	PERCENT
346-15 (N)	233	177	121	28	0	0	559	8.20
16 - 45 (NNE)	64	96	85	36	0	0	281	4.12
46 75 (ENE)	77	165	122	18	2	0	384	5.63
76 - 105 (E)	65	265	334	82	37	1	784	11.50
106 - 135 (ESE)	82	261	324	48	1	0	716	10.50
136 - 165 (SSE)	69	264	317	151	7	0	808	11.86
166 - 195 (S)	84	155	201	148	19	0	607	8.91
196 - 225 (SSW)	121	150	106	50	4	6	437	6.41
226 - 255 (WSW)	113	129	92	53	27	14	428	6.28
256 - 285 (W)	146	117	59	50	22	11	405	5.94
286 - 315 (WNW)	185	144	120	158	9	1	617	9.05
316 - 345 (NNW)	254	251	136	11	0	0	652	9.57
CALM	138						138	2.03
TOTAL	1631	2174	2017	833	128	33	6816	
PERCENT	23.93	31.90	29.59	12.22	1.88	0.48		100.00

TABLE 4

SITE #21 - ROYAL PALM BEACH  
WIND DIRECTION AND SPEED (MI/HR) OCCURRENCES  
1983

DIRECTION	1-3	4-6	7-10	11-16	17-21	> 22	TOTAL	PERCENT
346-15 (N)	83	3	0	0	0		86	1.02
16 - 45 (NNE)	128	9	0	0	0		137	1.63
46 75 (ENE)	224	26	0	0	0		250	2.97
76 - 105 (E)	653	306	48	2	0		1009	11.99
106 - 135 (ESE)	432	303	84	0	0		819	9.73
136 - 165 (SSE)	277	171	93	9	0		550	6.53
166 - 195 (S)	195	59	23	7	0		284	3.37
196 - 225 (SSW)	150	68	24	7	0		249	2.96
226 - 255 (WSW)	178	46	34	4	1		263	3.13
256 - 285 (W)	217	71	53	19	0		360	4.28
286 - 315 (WNW)	276	50	16	0	0		342	4.06
316 - 345 (NNW)	497	29	0	0	0		526	6.25
CALM	3542						3542	42.08
TOTAL	6852	1141	375	48	1		8417	
PERCENT	81.41	13.56	4.45	0.57	0.01			100.00



TABLE 5

SITE #16 - 20 MILE BEND  
WIND DIRECTION AND SPEED (MI/HR) OCCURRENCES  
1983

DIRECTION	1-3	4-6	7-10	11-16	17-21	> 22	TOTAL	PERCENT
346-15 (N)	109	39	15	2	0	0	165	3.06
16 - 45 (NNE)	174	63	24	0	0	0	261	4.83
46 75 (ENE)	202	35	12	0	0	0	249	4.61
76 - 105 (E)	200	112	57	18	1	0	388	7.19
106 - 135 (ESE)	346	161	59	10	0	0	576	10.67
136 - 165 (SSE)	356	174	112	22	0	0	664	12.30
166 - 195 (S)	107	86	67	13	0	0	273	5.05
196 - 225 (SSW)	160	54	25	17	0	0	256	4.74
226 - 255 (WSW)	220	37	30	6	0	0	293	5.43
256 - 285 (W)	122	39	25	24	7	1	218	4.04
286 - 315 (WNW)	234	97	90	63	6	0	490	9.07
316 - 345 (NNW)	221	60	24	7	0	0	312	5.78
CALM	1254						1254	23.23
TOTAL	3705	957	540	182	14	1	5399	
PERCENT	68.62	17.73	10.00	3.37	0.26	0.02		100.00

III  
TECHNICAL STUDIES

INTRODUCTION

The ambient air monitoring program in Palm Beach County during the year 1983 consisted of the following:

Total Suspended Particulate - 10 Sites

Total Gravimetric

Continuous Gaseous Monitoring - 4 Sites

Site 1

Nitrogen Dioxide

Carbon Monoxide

Site 22

Sulfur Dioxide

Sites 16 & 21

Ozone

Continuous Meteorological Monitoring - 4 Sites

Site 1

Wind Speed

Wind Direction

Temperature

Relative Humidity

Sites 16 & 21

Wind Speed

Wind Direction

Site 15

Temperature Inversion Monitoring

Microscopic Morphology

All criteria pollutant data collected is reported monthly to the Department of Environmental Regulation and to the Environmental Protection Agency for inclusion in air quality data banks (SAROAD). Gaseous pollutant levels and meteorological conditions from site #1 are relayed by data line to the agency's office data acquisition system. Instantaneous levels of these parameters are available at all times. One, three, eight and twenty-four hour averages are calculated and recorded. A Sum X Corporation Data Acquisition System was installed linking site #21 and the main office in October 1983. This site monitors for O<sub>3</sub>, wind speed, wind direction and trailer interior temperature.

Locations of the monitoring sites are shown in Figure 1. Table 6 gives site identification numbers, addresses, and parameters measured for all monitoring sites within the County. Table 7 relates measured air quality within Palm Beach County for the year 1983 to the Federal and State Ambient Air Quality Standards.

#### PERMANENT MONITORING NETWORK

The original monitoring network for suspended particulate sites, one through eight, was established in 1969 (Site 1A excluded). The Military Trail intercept line, sites nine through twelve, was added in 1972. Six of these twelve stations have been maintained as suspended particulate monitoring sites through the report period.

Periodic automated monitoring of sulfur dioxide, nitrogen dioxide and total oxidants (sites 1 through 8) was begun in June of 1971. Total hydrocarbon monitoring was begun in May of



1972 and discontinued in August of 1981. The automated gaseous and meteorological monitoring equipment were installed in the West Palm Beach monitoring station (Site 1) in November of 1972. Original Technicon Monitoring equipment for nitrogen dioxide, sulfur dioxide and total oxidants was replaced during the third quarter of 1973. Total oxidant monitoring was replaced by ozone monitoring at this time. Site 1 was maintained as the central monitoring station for gaseous pollutants, T.S.P., and meteorological parameters until March 1978, when the ozone monitor was relocated in Royal Palm Beach (Site 21). A second rural ozone monitor was placed on line in January 1980 at the South Florida Water Management Pump Station (Site 16) in accord with the National Air Monitoring Stations (NAMS) network. Relocation of the SO<sub>2</sub> monitoring site to Riviera Beach (Site 22), in July 1980, completed the State and Local Air Monitoring Stations (SLAMS). Network design is summarized in Table 8.

A special study of sulfur dioxide levels and meteorological parameters was conducted in Belle Glade (Site 8), from September of 1972 until May 1978.

Two manual stations for the measurement of nitrogen dioxide were established in November 1973 and maintained until October of 1978 as required by the State Implementation Plan. Manual sulfur dioxide stations have been operated periodically as part of special study projects.

#### PARTICULATE MONITORING

Methodology: Standard High volume samplers and shelters are located at each of the ten sites. The Belle Glade site,

#19, was relocated to Site #23 on November 4, 1983. Samples are collected and handled in accordance with Referenced Method for the Determination of Suspended Particulates in the Atmosphere (High Volume Method), Federal Register, Vol. 36, No. 84 - Friday, April 30, 1971. Sampling time is twenty-four hours, running from midnight to midnight, for each sampling date. The standard six day schedule as recommended by EPA is followed.

Tabulated results for suspended particulate for the year 1983 are presented in Tables 9 and 10. Figure 2 presents the range of probable logarithmic values for suspended particulate at all ten stations for the years 1982 and 1983. A historical summation for total suspended particulate measurements from 1969 thru 1983 is presented in Table 11.

#### GASEOUS MONITORING

Maximum ambient air concentrations for gaseous sampling in Palm Beach County for the period 1970-1983 are presented in Table 12.

#### NITROGEN DIOXIDE (SLAMS)

Continuous automatic monitoring for this pollutant is carried on at Site 1. A MEC Model 1200 NO-NO<sub>x</sub> (McMillan Electronics Corporation) Chemiluminescence analyzer was in use from November 15, 1973 until December 27, 1977. At this time it was replaced by Monitor Laboratory Model 8440. Table 13 presents monthly and annual sampling time, arithmetic means and twenty-four hour maximum concentrations. Table 14 includes quarterly and annual, one, eight and twenty-four hour concentration maximums, annual arithmetic means and the frequency distribution of

ranges of pollutant levels recorded. Values recorded placed the measured concentrations of this pollutant well below those of the Ambient Air Quality Standards. D.E.R. was unable to make NO<sub>2</sub> accuracy audits during the second half of the year. Consequently, accuracy data for this pollutant is available for only the first half of the year. Precision and completeness for the year are as follows:

Accuracy	
Concentration Range	Accuracy Range
0.08 - 0.10 ppm	-73 to 33%
0.15 - 0.10 ppm	-50 to 17%
0.40 - 0.45 ppm	-48 to 14%
Precision:	-21 to 12%
Completeness:	83%

#### CARBON MONOXIDE (SLAMS)

This pollutant is monitored continuously at Site 1. A Mine Safety Appliances, Model 200 nondispersive infrared spectrophotometric automatic analyzer, in service since January 1971, was retired and replaced by a Model 202-S on October 6, 1977. Table 15 presents a monthly record of sampling hours, one and eight hour maximums, and relates concentration maximums to the Ambient Air Quality Standard. There were no recorded values in excess of the one hour or eight hour standards during the reported period.

Table 16 presents quarterly, one and eight hour maximum values and frequency distribution of all recorded pollutant



levels. Accuracy, precision and completeness for the year are as follows:

Accuracy	
Concentration Range	Accuracy Range
3 - 8 ppm	-16 to 13%
15 - 20 ppm	- 4 to 5%
40 - 45 ppm	- 4 to 5%
Precision: -16 to 8%	
Completeness: 71%	

#### OZONE (NAMS)

This pollutant was monitored continuously at Site 1 from September 6, 1973 thru 1978. The instrumentation is a MEC Model 1100 (McMillan Electronic Corporation) Chemiluminescence analyzer. In 1978 the instrument was modified by the manufacturer to EPA designated reference method status. The monitor was re-located to Royal Palm Beach, Site 21, and placed in service there March 1, 1979.

In January, 1980 a Monitor Labs Model 8410 ozone monitor, went on line at the South Florida Water Management Pump Station (Site 16). This completed the EPA mandated NAMS ozone monitoring network for Palm Beach County. On December 19, 1983 these two monitors were reassigned, the Monitor Lab was transferred to Royal Palm Beach and the McMillan to the South Florida Pump Station.

The Dasibi Model #1003 PC, which had functioned as the calibration system since December, 1978 was reassigned to trans-

fer calibrator status in 1980 in conjunction with a Columbia Scientific Photocal 3000 primary standard.

Tables 17 and 19 present a monthly record of hours sampled and one hour maximum recorded at each site. There were no values recorded which exceed the Federal Ambient Air Quality Standard for this pollutant.

One, eight and twenty-four hour maximum and the frequency distribution for all pollutant levels recorded are presented on a quarterly basis in Tables 18 and 20.

Tables 21 and 22 give the design values or expected maximum hourly averages for these sites based on yearly data and on last three years data Wiebull distributions. No credit is given for "seasonal free days" in these calculations. Accuracy precision and completeness for the year are as follows:

Accuracy	
Concentration Range	Accuracy Range
0.08 - 0.10 ppm	-3 to 7%
0.15 - 0.10 ppm	-4 to 7%
0.40 - 0.45 ppm	-8 to 8%

Precision: -8 to 14%

Completeness: 82%

#### SULFUR DIOXIDE (SLAMS)

Monitoring instrumentation is a Monitor Labs Model 8850 Fluorescent monitor which replaced the Beckman 904-A Sulfur Dioxide analyzer, based on coulometric titration.

Site 22, located in Riviera Beach was established in July, 1980 to monitor sulfur dioxide levels in the coastal region of the County. Table 23 reports a monthly history of hours sampled, one, three, and twenty-four hour maximums, and the number of violations for each of the related standards. Table 24 presents quarterly and annual maximum values and a frequency distribution of values recorded. Calculation of an arithmetic mean for this pollutant is not feasible because ninety-nine percent of the values are below the limit of sensitivity for the instrument. Precision, accuracy and completeness for the year are as follows:

Accuracy

Concentration Range	Accuracy Range
0.03 - 0.08 ppm	2 to 6%
0.15 - 0.20 ppm	-2 to 14%
0.40 - 0.45 ppm	0 to 11%

Precision: -15 to 7%

Completeness: 95%



TABLE 6

## MONITORING SITE LOCATIONS

SITE NO.	ADDRESS	UTM ZONE 17	MONITORING CAPABILITY
1	West Palm Beach Water Treatment Plant First Street and Tamarind Avenue West Palm Beach, Florida	2955030N 0593232E	CO, NO <sub>2</sub>  Meteorology
1A	Palm Beach County Health Department 901 Evernia Street West Palm Beach, Florida	2955030N 0593232E	Susp. Part.
2	Tequesta Water Department 357 Tequesta Drive Tequesta, Florida	2982018N 0589963E	Discontinued December 27, 1977
3	North Palm Beach Water Treatment Plant 603 Anchorage Drive North Palm Beach, Florida	2965817N 0592780E	Susp. Part.
4	Lake Worth Water Treatment Plant 301-303 College Street Lake Worth, Florida	2943537N 0592793E	Susp. Part.
5	Delray Beach Water Treatment Plant 202 N. W. 1st Avenue Delray Beach, Florida	2927488N 0592195E	Susp. Part.
6	Boca Raton Fire Station #1 1151 N. Federal Highway Boca Raton, Florida	2915768N 05913137E	Susp. Part.
7	Royal Palm Beach Golf Course Royal Palm Beach Boulevard Royal Palm Beach, Florida	2951437N 0578767E	Discontinued March 27, 1978
8	Belle Glade Water Treatment Plant 1016 West Canal Street Belle Glade, Florida	2953082N 0533160E	Discontinued October 29, 1978
9	Grammercy Park Water Treatment Plant Park Avenue Grammercy Park, Florida	2960537N 0587329E	Discontinued February 1, 1977

TABLE 6

## MONITORING SITE LOCATIONS (CONT'D)

SITE NO.	ADDRESS	UTM ZONE 17	MONITORING CAPABILITY
10	Southwest Fire Department 1180 S. Military Trail West Palm Beach, Florida	2949018N 0588207E	Susp. Part.
11	St. Vincent DePaul Seminary S. Military Trail Boynton Beach, Florida	2932890N 0586927E	Discontinued July 17, 1976
12	College of Boca Raton S. Military Trail Boca Raton, Florida	2918354N 0587320E	Susp. Part.
13	NO <sub>x</sub> SIP Site N8 Florida Atlantic University Boca Raton, Florida	2917000N 0589500E	Discontinued October 1, 1978
14	NO <sub>x</sub> SIP Site N9 Palm Beach Mall Palm Beach Lakes Boulevard West Palm Beach, Florida	2956000N 0590700E	Discontinued October 1, 1978
15	Division of Forestry Lat. 26° 41'N, Long 80° 16'E Loxahatchee, Florida		Temperature Inversion
16	South Florida Water Management Pump Station Twenty Mile Bend State Road 80	2951402N 0562879E	O <sub>3</sub> Susp. Part. Meteorology
17	Lake Harbor Water Treatment Plant Lake Harbor, Florida	2952230N 0518600E	Discontinued December 27, 1977
18	Pahokee Health Department 1759 E. Main Street Pahokee, Florida	2967222N 0533760E	Discontinued November 1, 1978
19	Belle Glade Fire Station 22 W. Avenue "A" Belle Glade, Florida	2951420N 0532900E	Discontinued November 4, 1983
20	Pahokee Sewage Treatment Plant 1050 McClure Road Pahokee, Florida	2964200N 0532300E	Susp. Part.

TABLE 6

## MONITORING SITE LOCATIONS (CONT'D)

SITE NO.	ADDRESS	UTM ZONE 17	MONITORING CAPABILITY
21	Royal Palm Beach R. V. Area 10999 Okeechobee Boulevard Royal Palm Beach, Florida	2954150N 0578100E	O <sub>3</sub> , Meteorology
22	Palm Beach County Health Department Warehouse 2030 Avenue "L" Riviera Beach, Florida	296235N 059248E	SO <sub>2</sub>
23	Belle Glade Health Department 1024 N. W. Avenue "D" Belle Glade, Florida	2953082N 0533160E	Susp. Part.






Figure 1 Ambient Air Monitoring Network

TABLE 7

## AMBIENT AIR QUALITY STANDARDS

POLLUTANT	FEDERAL PRIMARY	FEDERAL SECONDARY	STATE	MEASURED LEVELS PALM BEACH COUNTY PPM see table #			
				Site 1	Site 16	Site 21	Site 22
Suspended Particulates							
Annual Geo. Mean	75 ug/m <sup>3</sup>	60 ug/m <sup>3</sup>	60 ug/m <sup>3</sup>				
Maximum 24 hr. Conc. (2)	260 ug/m <sup>3</sup>	150 ug/m <sup>3</sup>	150 ug/m <sup>3</sup>				
Sulfur Oxides							
Annual Arith. Mean	80 ug/m <sup>3</sup> (0.03 ppm)		60 ug/m <sup>3</sup> (0.02 ppm)				0.0005
Maximum 24 hr. Conc.	365 ug/m <sup>3</sup> (0.14 ppm)		260 ug/m <sup>3</sup> (0.1 ppm)				0.015
Maximum 3 hr. Conc. (2)		1,300 ug/m <sup>3</sup> (0.5 ppm)	1,300 ug/m <sup>3</sup> (0.5 ppm)				0.025
Carbon Monoxide							
Maximum 8 hr. Conc. (2)	10 mg/m <sup>3</sup> (9 ppm)	Same as Federal Primary	Same as Federal Primary	6.5			
Maximum 1 hr. Conc.	40 mg/m <sup>3</sup> (35 ppm)			8.9			
Ozone							
Daily Maximum 1 hr. Conc. (1)	235 ug/m <sup>3</sup> (0.12 ppm)	Same as Federal Primary	Same as Federal Primary		0.087	0.092	
Nitrogen Oxides							
Annual Arith. Mean	100 ug/m <sup>3</sup> (0.05 ppm)	Same as Federal Primary	Same as Federal Primary	0.010			

1. The standard is attained when the expected number of days per calendar year with maximum hourly average concentrations above 0.12 ppm is equal to or less than 1.
2. Concentration limits not to be exceeded more than once per year.

TABLE 8

PALM BEACH COUNTY  
NETWORK DESCRIPTION  
NAMS, SLAMS AND SPECIAL PURPOSE

Urban Area	Site Address	Network	Saroad Site #	Sampler	Analysis	Pollutant	Mon.Obj.	Spatial Scale	Ope. Schedule	Implem.
Belle Glade	1024 NW Ave. "D"	SLAMS	0240-003-G01	Hi Vol	Gravimetric	TSP	Max.Conc.	Neighborhood	6 day	Operational
Boca Raton	1151 N. Fed. Hwy	SLAMS	0280-001-G01	Hi Vol	Gravimetric	TSP	Pop.Exp.	Neighborhood	6 day	Operational
Boca Raton	S. Military Trail	SLAMS	0280-002-G01	Hi Vol	Gravimetric	TSP	Pop.Exp.	Neighborhood	6 day	Operational
Delray Beach	202 NW 1st Ave.	SLAMS	1000-002-G01	Hi Vol	Gravimetric	TSP	Max.Conc.	Neighborhood	6 day	Operational
Lake Worth	301 College St.	SLAMS	2220-001-G01	Hi Vol	Gravimetric	TSP	Max.Conc.	Neighborhood	6 day	Operational
N.Palm Beach	603 Anchorage Dr.	SLAMS	3060-001-G01	Hi Vol	Gravimetric	TSP	Pop.Exp.	Neighborhood	6 day	Operational
W.Palm Beach	1108 S. Military Trail	SLAMS	3420-005-G01	Hi Vol	Gravimetric	TSP	Pop.Exp.	Neighborhood	6 day	Operational
Rural	20 Mile Bend Rd.	NAMS	3120-006-G03	Monitor Lab 8410	Chemilum	O <sub>3</sub>	Max.Conc.	Urban	cont.	Operational
Rural	10999 Okeechobee Blvd.	NAMS	3420-007-G01	MEC 1100-1	Chemilum	O <sub>3</sub> (EE)	Pop.Exp.	Neighborhood	cont.	Operational
Riviera Beach	2030 Ave. "L"	SLAMS	3840-003-G02	Monitor Lab 8850	Fluorescent	SO <sub>2</sub>	Max.Conc.	Neighborhood	cont.	Operational
W.Palm Beach	1st St. & Tamarind Ave	SLAMS	4760-001-G01	Monitor Lab 8440	Chemilum	NO <sub>2</sub>	Max.Conc.	Neighborhood	cont.	Operational
W.Palm Beach	901 Evernia St.	SLAMS	4760-003-G01	Hi Vol	Gravimetric	TSP	Max.Conc.	Neighborhood	6 day	Operational
W.Palm Beach	1st St. & Tamarind Ave	SLAMS	4760-001-G01	MSA 202-S	Nondispersive Infrared	CO	Max.Conc.	Neighborhood	cont.	Operational
Rural	Twenty Mile Bend	S.P.	3420-006-G01	Hi Vol	Gravimetric	TSP	Max.Conc.	Neighborhood	6 day	Operational
Pahokee	1050 McClure Rd.	S.P.	3340-001-G01	Hi Vol	Gravimetric	TSP	Max.Conc.	Neighborhood	6 day	Operational

Note: On December 19, 1984 the O<sub>3</sub> monitors were changed. The Monitor Lab 8410 was placed at the 10999 Okeechobee Blvd. site and the MEC 1100-1 was placed at the 20 Mile Bend site.

TABLE 9

## SUSPENDED PARTICULATE MATTER, 1983

GEOMETRIC MEAN ug/m <sup>3</sup>						Geo. Std. Dev.	CONCENTRATION ug/m <sup>3</sup>				% above or below Annual Standard (60 ug/m <sup>3</sup> )	No. of Samples Above Daily Standard (150 ug/m <sup>3</sup> )
Site No.	1st Qtr.	2nd Qtr.	3rd Qtr.	4th Qtr.	Annual		Arith. Mean	Max.	2nd Max.	Min.		
1A	42.2	31.4	32.8	26.6	32.9	1.57	36.8	124	103	11	-45.17	-0-
3	30.3	24.6	27.5	20.9	25.6	1.56	28.4	117	101	11	-57.33	-0-
4	45.9	35.2	35.0	26.4	35.0	1.54	38.7	130	98	13	-41.67	-0-
5	41.7	35.2	35.9	25.8	34.1	1.80	37.2	126	107	14	-43.17	-0-
6	37.9	32.9	35.7	27.7	33.4	1.48	36.3	134	90	14	-44.33	-0-
10	50.0	37.0	42.2	30.6	39.5	1.50	42.9	122	104	16	-34.17	-0-
12	39.4	27.1	26.7	21.6	27.2	1.57	30.4	116	82	12	-54.67	-0-
16	32.2	25.0	24.8	20.4	25.2	1.54	27.8	73	69	9	-58.00	-0-
19	53.1	42.8	39.6	38.1	43.1	1.42	45.8	102	84	23	-28.17	-0-
20	38.6	41.8	42.0	31.7	38.1	1.41	40.5	100	86	19	-36.50	-0-



TABLE 10

TOTAL SUSPENDED PARTICULATES - 1983  
SAMPLING DAYS

SITE NO.	NUMBER OF SAMPLES												
	J	F	M	A	M	J	J	A	S	O	N	D	TOTAL
1A	5	4	6	5	5	5	5	5	5	5	5	4	59
3	5	4	6	4	5	5	4	5	5	5	5	5	58
4	5	4	6	5	5	5	5	5	5	5	5	5	60
5	4	4	6	5	4	5	3	5	5	4	5	5	55
6	5	4	6	4	5	5	5	5	5	4	5	5	58
10	5	4	6	5	4	5	5	5	5	5	5	4	58
12	2	3	5	5	5	5	5	5	5	5	5	5	55
16	5	4	6	5	5	5	5	5	5	5	5	5	60
19	5	4	6	5	5	5	5	5	5	5	5	4	59
20	4	3	6	5	5	5	4	4	3	4	5	5	53

FIGURE 2






TABLE 11  
TOTAL SUSPENDED PARTICULATE ug/m<sup>3</sup>  
1969-1983

Site No.	Parameter	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983
1/1A	Maximum	109.3	115.7	121.1	133.6	101.9	96.4	81.5	106.2	172	92	88	78	119	72	124
	Minimum	9.6	13.9	15.2	15.2	10.8	19.6	20.0	15.8	17	18	18	22	20	11	11
	Arith. Mean	48.7	40.9	58.7	49.9	40.5	40.9	44.7	37.4	42.0	41.3	41.0	43.9	49.4	35.3	36.8
	Geo. Mean	43.9	39.5	53.4	45.9	38.0	38.8	42.4	35.3	38.3	38.2	38.5	42.2	45.8	33.2	32.9
	Geo. Std. Dev.	1.56	1.42	1.19	1.49	1.49	1.38	1.39	1.41	1.50	1.47	1.44	1.34	1.48	1.45	1.57
2	Maximum	71.2	74.3	122.3	112.3	85.4	104.0	77.7	63.1	74						
	Minimum	7.3	13.3	10.7	12.2	5.3	19.6	11.9	14.0	12						
	Arith. Mean	26.9	30.4	34.6	33.0	33.6	32.4	36.4	35.5	33.6						
	Geo. Mean	24.6	28.3	30.8	30.3	31.4	30.6	34.1	33.4	30.9						
	Geo. Std. Dev.	1.51	1.45	1.61	1.48	1.49	1.36	1.45	1.44	1.50						
3	Maximum	71.8	82.3	167.5	94.8	133.2	132.7	91.8	67.8	62	65	85	90	115	59	117
	Minimum	7.7	1.3	0.4	12.8	16.6	16.1	14.6	10.3	11	12	14	20	17	12	11
	Arith. Mean	32.2	31.7	40.6	37.0	38.2	35.8	38.3	31.4	30.4	32.1	37.8	41.5	42.6	28.4	28.4
	Geo. Mean	29.5	28.4	30.7	33.7	35.3	32.2	34.5	29.1	28.4	30.2	35.3	39.2	39.0	26.4	25.6
	Geo. Std. Dev.	1.63	1.76	2.93	1.49	1.47	1.54	1.61	1.50	1.45	1.42	1.47	1.41	1.53	1.49	1.56
4	Maximum	351.9	224.8	95.6	89.8	85.6	196.8	435.3	81.0	84	85	101	90	123	121	130
	Minimum	7.3	8.0	10.2	12.9	13.3	18.3	19.6	12.7	14	17	20	9	25	14	13
	Arith. Mean	32.9	30.9	37.2	34.8	37.7	45.2	57.1	38.2	41.0	44.6	44.7	47.8	49.0	38.2	38.7
	Geo. Mean	26.4	28.2	31.7	32.2	35.3	38.8	47.9	35.8	39.1	42.1	42.0	45.0	45.3	35.1	35.0
	Geo. Std. Dev.	1.78	1.47	1.85	1.49	1.39	1.67	1.64	1.44	1.38	1.41	1.43	1.46	1.48	1.53	1.54
5	Maximum	164.9	76.7	142.4	108.0	92.9	81.9	83.5	61.1	81	99	102	82	122	76	126
	Minimum	13.3	8.3	12.2	15.9	10.9	13.8	22.0	13.8	18	17	19	26	21	16	14
	Arith. Mean	40.1	36.2	36.4	38.5	40.0	34.8	42.0	35.8	39.1	37.6	40.4	42.7	46.2	35.4	37.2
	Geo. Mean	38.8	33.6	32.0	35.4	37.6	32.2	39.5	34.0	37.0	35.0	37.6	41.1	42.5	33.2	34.1
	Geo. Std. Dev.	1.47	1.49	1.64	1.49	1.46	1.53	1.41	1.40	1.41	1.45	1.47	1.33	1.5	1.43	1.80

TABLE 11  
TOTAL SUSPENDED PARTICULATE ug/m<sup>3</sup>  
1969-1983

Site No.	Parameter	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983
6	Maximum	83.1	80.1	327.9	275.3	106.5	92.4	114.8	62.8	79	107	124	94	131	70	134
	Minimum	9.5	9.6	13.3	17.0	13.6	20.9	22.4	16.4	15	18	19	26	25	15	14
	Arith. Mean	36.4	33.5	49.1	44.9	43.1	41.6	45.4	35.6	39.3	42.2	47.3	47.0	48.6	27.8	36.3
	Geo. Mean	32.9	31.0	41.1	39.9	40.2	38.4	42.7	33.8	37.0	39.3	43.8	44.8	45.7	32.6	33.4
	Geo. Std. Dev.	1.60	1.48	2.09	1.56	1.46	1.26	1.43	1.38	1.43	1.45	1.48	1.37	1.42	1.47	1.48
7	Maximum	52.5	71.7	131.5	102.0	65.5	98.3	70.5	55.2	64	36					
	Minimum	7.2	2.1	1.6	7.0	9.1	5.4	11.4	6.2	9	17					
	Arith. Mean	23.6	25.8	30.7	31.8	28.1	25.6	33.0	23.1	24.3						
	Geo. Mean	21.5	23.3	24.4	28.3	26.2	22.3	30.4	21.0	22.5						
	Geo. Std. Dev.	1.57	1.59	2.13	1.37	1.45	1.66	1.52	1.55	1.49						
8	Maximum	175.7	273.9	222.7	173.3	151.0	210.9	199.4	125.2	149	143					
	Minimum	12.7	14.5	12.6	19.8	20.3	22.8	10.7	12.6	17	22					
	Arith. Mean	53.8	54.6	61.4	58.6	59.8	59.8	62.4	61.6	59.0	58.8					
	Geo. Mean	46.0	47.1	53.1	52.3	54.0	54.2	56.7	56.3	54.6	53.1					
	Geo. Std. Dev.	1.76	1.70	1.64	1.60	1.61	1.57	1.56	1.56	1.49	1.57					
9	Maximum				74.50	145.3	81.2	65.3	59.1	33						
	Minimum				13.30	11.7	11.3	16.1	9.6	20						
	Arith. Mean				31.2	33.2	29.9	34.1	28.2							
	Geo. Mean				28.7	30.7	27.0	32.2	26.3							
	Geo. Std. Dev.				1.42	1.45	1.50	1.39	1.46							
10	Maximum				94.80	109.0	113.0	81.7	101.6	98	77	80	87	122	72	122
	Minimum				18.30	19.0	21.0	23.4	12.8	19	21	18	23	19	17	16
	Arith. Mean				44.4	45.3	43.0	47.2	42.8	41.4	46.0	46.2	51.2	53.1	38.2	42.9
	Geo. Mean				41.6	42.5	39.0	45.4	40.5	39.2	43.6	43.3	48.9	48.4	36.2	39.5
	Geo. Std. Dev.				1.43	1.43	1.54	1.35	1.40	1.40	1.41	1.45	1.33	1.54	1.40	1.50



TABLE 11

TOTAL SUSPENDED PARTICULATE ug/m<sup>3</sup>  
1969-1983

Site No.	Parameter	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983
11	Maximum				69.9	77.8	134.3	299.9	60.8							
	Minimum				11.9	11.5	13.5	13.5	10.9							
	Arith. Mean				32.1	30.8	34.4	44.9								
	Geo. Mean				29.2	28.9	29.4	37.7								
	Geo. Std. Dev.				1.51	1.53	1.70	1.67								
12	Maximum				68.1	79.6	200.7	64.7	53.7	66	69	85	107	122	62	116
	Minimum				11.2	11.0	12.8	18.2	7.0	14	14	13	14	18	10	12
	Arith. Mean				29.6	31.7	34.9	34.2	26.9	29.8	29.0	35.0	38.5	40.8	27.3	30.4
	Geo. Mean				26.9	29.6	29.2	32.4	25.1	27.9	27.2	31.7	35.8	37.0	25.0	27.2
	Geo. Std. Dev.				1.52	1.43	1.70	1.39	1.47	1.43	1.43	1.56	1.54	1.55	1.52	1.57
16	Maximum								130.1	76	136	87	68	96	128	73
	Minimum								14.6	12	14	14	19	16	9	9
	Arith. Mean								35.0	30.9	31.5	37.3	34.2	43.4	26.5	27.8
	Geo. Mean								31.0	38.2	28.7	33.8	32.1	38.4	23.5	25.2
	Geo. Std. Dev.								1.60	1.52	1.50	1.57	1.44	1.66	1.60	1.54
17	Maximum									69						
	Minimum									10						
	Arith. Mean									34.6						
	Geo. Mean									31.6						
	Geo. Std. Dev.									1.53						
18/20	Maximum									63	76					
	Minimum									9	16					
	Arith. Mean										30.8					
	Geo. Mean										28.8					
	Geo. Std. Dev.										1.44					

TABLE 11

TOTAL SUSPENDED PARTICULATE  $\mu\text{g}/\text{m}^3$   
1969-1983

Site No.	Parameter	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983
19/23	Maximum										121	121	110	166	87	102
	Minimum										20	16	31	25	16	23
	Arith. Mean										52.9	57.5	58.9	62.7	47.8	45.8
	Geo. Mean										49.6	53.9	56.5	56.6	45.1	43.1
	Geo. Std. Dev.										1.45	1.45	1.34	1.57	1.41	1.41
20	Maximum											122	164	117	85	100
	Minimum											21	16	21	12	19
	Arith. Mean											50.6	52.0	55.6	41.8	40.5
	Geo. Mean											46.4	47.9	50.6	38.6	38.1
	Geo. Std. Dev.											1.52	1.48	1.56	1.52	1.41

TABLE 12

GASEOUS SAMPLING  
MAXIMUM AMBIENT AIR CONCENTRATIONS, PPM  
1970-1983

Site No.	Sampling Dates	Sulfur Dioxide			Total Oxidants		Ozone	
		1 Hour	3 Hour	24 Hour	1 Hour	8 Hour	1 Hour	8 Hour
1	07/17-07/31/70	0.094	0.038	0.007	0.114	0.073	-	-
	04/12-04/27/71	0.044	0.028	> 0.006	> 0.188	> 0.130	-	-
	07/16-07/30/71	0.035	0.012	0.002	0.032	0.026	-	-
	11/14-12/31/72	0.023	0.021	0.003	> 0.187	> 0.040	-	-
	01/01-11/14/73	0.042	0.034	0.004	-	-	-	-
	11/14-12/31/73	0.015	0.008	0.001	-	-	-	-
	01/01-11/01/73				0.155	0.063		
	09/06-12/31/73						> 0.100	> 0.071
	01/01-12/31/74	0.075	0.052	0.023			0.077	0.055
	01/01-12/31/75	0.062	0.025	0.008	-	-	0.104	0.077
	01/01-12/31/76	0.055	0.034	0.021	-	-	0.148	0.122
	01/01-12/31/77	0.019	0.015	0.009	-	-	0.106	0.088
	01/01-12/31/78				-	-	0.075	0.066
	04/01-06/30/78	0.030	0.022	0.008	-	-	-	-
	01/01-10/22/79	0.024	0.017	0.016	-	-	-	-
2	06/16-07/02/70	0.035	0.026	0.010	0.104	0.093	-	-
	05/11-05/25/71	0.191	0.142	0.028	0.010	0.0004	-	-
	08/13-08/27/71	0.033	0.015	0.003	0.016	0.018	-	-
3	07/02-07/17/70	0.196	0.028	0.028	0.176	0.086	-	-
	04/27-05/11/71	> 0.500	0.324	0.060	0.111	0.055	-	-
	07/30-08/13/71	0.064	0.035	0.005	0.007	0.001	-	-
	05/18-06/30/72	0.053	0.032	0.006	0.116	0.071	-	-
	09/03-12/31/76	-	-	0.004	-	-	-	-
	01/01-12/31/77	-	-	0.004	-	-	-	-

TABLE 12

GASEOUS SAMPLING  
 MAXIMUM AMBIENT AIR CONCENTRATIONS, PPM  
 1970-1983  
 (continued)

Site No.	Sampling Dates	Sulfur Dioxide			Total Oxidants		Ozone	
		1 Hour	3 Hour	24 Hour	1 Hour	8 Hour	1 Hour	8 Hour
4	07/31-08/14/70	0.031	0.024	0.010	0.129	0.089	-	-
	03/26-04/12/71	0.044	0.034	0.012	0.110	0.106	-	-
	09/23-10/04/71	0.080	0.035	0.006	0.056	0.048	-	-
	11/10-11/19/71	0.000	0.000	0.000	0.078	0.073	-	-
5	09/04-09/18/70	0.069	0.029	0.003	0.092	0.066		
	03/12-03/26/71	0.060	0.028	0.005	0.013	0.086		
	10/19-11/01/71	0.006	0.002	0.0003	0.136	0.101		
6	08/21-09/04/70	0.076	0.048	0.013	0.048	0.037		
	01/27-02/12/71	0.132	0.098	0.017	0.110	0.095		
	12/29-01/12/72	0.068	0.034	0.006	0.000	0.000		
	07/05-08/01/72	0.015	0.012	0.003	0.050	0.027		
7	09/28-10/12/70	0.106	0.048	0.006	0.076	0.068		
	02/26-03/12/71	0.026	0.008	➤ 0.003	0.110	0.093		
	11/19-12/10/71	0.015	0.006	0.001	0.038	0.016		
8	10/12-10/26/70	0.000	0.000	0.000	0.078	0.061		
	02/12-02/26/71	➤ 0.288	0.125	➤ 0.030	0.103	0.076		
	02/10-12/29/71	➤ 0.267	0.217	➤ 0.039	0.012	0.006		
	09/21/72-05/01/73	0.068	0.028	0.007				
	12/01/72-05/18/73	0.176	0.098	0.044				
	12/18-12/31/73	0.153	0.113	0.025				
	01/01-09/29/74	0.065	0.031	0.004				
	07/08-12/31/75	0.075	0.074	0.029				
	01/01-12/31/76	0.085	0.078	0.051				
	01/01-12/31/77	0.033	0.029	0.016				
	01/01-05/19/78	0.047	0.043	0.019				



TABLE 12

GASEOUS SAMPLING  
 MAXIMUM AMBIENT AIR CONCENTRATIONS, PPM  
 1970-1983  
 (continued)

Site No.	Sampling Dates	Sulfur Dioxide			Total Oxidants		Ozone	
		1 Hour	3 Hour	24 Hour	1 Hour	8 Hour	1 Hour	8 Hour
16	01/03-12/31/80						.098	.081
	01/01-12/31/81						.095	.074
	01/01-12/31/82						.080	.073
	01/01-12/31/83						.087	.071
17	09/27-12/31/76			0.003				
	01/01-12/31/77			0.003				
21	03/01-12/31/79						.079	.070
	01/01-12/31/80						.110	.080
	01/01-12/31/81						.103	.079
	01/01-12/31/82						.122	.073
	01/01-12/31/83						.092	.073
22	07/24-12/31/80	0.016	0.014	0.013				
	01/01-12/31/81	0.034	0.031	0.019				
	01/01-12/31/82	0.067	0.053	0.019				
	01/01-12/31/83	0.035	0.025	0.015				

TABLE 12

MAXIMUM AMBIENT AIR CONCENTRATION, PPM  
1970-1983

Site No.	Sampling Dates	Nitrogen Dioxide			Carbon Monoxide		Total Hydrocarbons	
		1 Hour	8 Hour	Ar./Mean	1 Hour	8 Hour	1 Hour	8 Hour
1	07/17-07/31/70	0.097	0.068	0.016				
	04/12-04/27/71	0.147	0.079	0.026				
	07/15-07/30/71	0.067	0.056	0.018	3.6	3.1		
	11/14-12/31/72	0.092	0.079	0.020	7.0	3.7	6.5	3.2
	01/01-11/15/73	0.060	0.047	0.007				
	01/01-12/31/73				8.9	6.3	5.5	3.3
	01/01-12/31/74	0.080	0.052	0.015	10.5	8.8	5.8	4.4
	01/01-12/31/75	0.125	0.083	0.015	8.6	5.0	5.2	3.0
	01/01-12/31/76	0.083	0.054	0.009	10.5	5.2	5.3	3.7
	01/01-12/31/77	0.071	0.044	0.017	11.8	8.5	5.2	3.6
	01/01-12/31/78	0.089	0.070	0.012	8.6	4.5		
	10/18-12/31/78						5.8	3.2
	01/01-11/28/79	0.078	0.056	0.016				
	01/01-11/08/79				7.8	3.1		
	01/01-12/31/79						8.3	2.9
	01/01-12/31/80	0.137	0.088	0.018	9.8	5.7	9.6	6.2
	01/01-12/31/81	0.152	0.107	0.012	13.3	5.8		
	01/01-8/14/81						8.4	3.5
	01/01-12/31/82	0.069	0.053	0.038	21.3	8.1		
	01/01-12/31/83	0.069	0.055	0.010	8.9	6.5		
2	06/16-07/02/70	0.044	0.032	0.010				
	05/11-05/25/71	0.054	0.040	0.013	2.2	0.3		
	08/12-08/27-71	0.073	0.060	0.013	0.0	0.0		

TABLE 12

MAXIMUM AMBIENT AIR CONCENTRATION, PPM  
1970-1983  
(continued)

Site No.	Sampling Dates	Nitrogen Dioxide			Carbon Monoxide		Total Hydrocarbons	
		1 Hour	8 Hour	Ar./Mean	1 Hour	8 Hour	1 Hour	8 Hour
3	07/02-07/17/70	0.084	0.060	0.010				
	04/27-05/11/71	0.096	0.066	0.017				
	07/30-08/13/71	0.083	0.069	0.018	3.2	0.9		
	05/18-06/30/72	0.088	0.059	0.010	0.0	0.0	3.2	2.2
	01/01-12/31/76			0.006				
	01/01-12/31/77			0.010				
	01/01-03/31/78			0.014				
4	07/31-08/14/70	0.097	0.068	0.016				
	03/26-04/12/71	0.118	0.107	0.018	2.1	0.3		
	09/23-10/14/71	0.059	0.041	0.018	0.0	0.0		
	11/10-11/19/71	0.124	0.101	0.020	0.0	0.0		
5	09/04-09/18/70	0.055	0.051	0.013				
	03/12-03/26/71	0.146	0.113	0.018	2.1	0.4		
	10/19-11/01/71	0.117	0.093	0.029				
6	08/21-09/04/70	0.064	0.048	0.015				
	01/27-02/12/71	0.200	0.187	0.047	9.6	4.2		
	12/29-01/12/71	0.079	0.069	0.022	2.6	0.4		
	07/05-08/01/72	0.065	0.055	0.011				
7	09/28-10/12/71	0.031	0.018	0.007				
	02/26-03/12/71	0.106	0.081	0.016	0.8	0.1		
	11/09-12/10/71	0.074	0.055	0.019	2.2	2.0		
8	10/12-10/26/70	0.118	0.067	0.017				
	02/12-02/26/71	0.152	0.091	0.022	5.2	3.0		
	12/10-12/29/71	0.076	0.048	0.024	0.0	0.0		

TABLE 12

MAXIMUM AMBIENT AIR CONCENTRATION, PPM  
1970-1983  
(continued)

Site No.	Sampling Dates	Nitrogen Dioxide			Carbon Monoxide		Total Hydrocarbons	
		1 Hour	8 Hour	Ar./Mean	1 Hour	8 Hour	1 Hour	8 Hour
13	11/14-12/31/73			0.003				
	01/01-12/31/74			0.004				
	01/01-12/31/75			0.008				
	01/01-12/31/76			0.005				
	01/01-12/31/77			0.008				
	01/01-12/31/78			0.010				
14	11/14-12/31/73			0.004				
	01/01-12/31/74			0.005				
	01/01-12/31/75			0.012				
	01/01-12/31/76			0.008				
	01/01-12/31/77			0.015				
	01/01-12/31/78			0.015				



TABLE 13

NITROGEN DIOXIDE DATA  
SITE NUMBER 1  
1983

Month	Hours	Arith. (a) Avg., ppm	Max. 24 Hr. Avg., ppm
J	515	.015	.032
F	527	.011	.019
M	345	.008	.021
A	701	.010	.019
M	548	.006	.011
J	688	.007	.018
J	583	.005	.012
A	590	.010	.022
S	709	.008	.020
O	726	.008	.020
N	581	.018	.031
D	737	.015	.035
Annual	7250	.010	.035

(a) Standard is 0.05 ppm, annual arithmetic mean.

TABLE 14

AMBIENT AIR, NITROGEN DIOXIDE, PPM  
SITE NUMBER 1  
1983

	Quarter 1	Quarter 2	Quarter 3	Quarter 4	Annual		
1 Hr. Maximum	.060	.061	.046	.069	.069		
8 Hr. Maximum	.048	.042	.039	.055	.055		
24 Hr. Maximum	.032	.019	.022	.035	.035		
Arith. Mean	.015	.010	.010	.018	.010		
Concentration Range			Hours			%	Cum. %
< .010	760	1356	1337	998	4451	61.39	61.39
.010 - .019	330	412	362	560	1664	22.95	84.34
.020 - .029	173	97	130	276	676	9.32	93.66
.030 - .039	84	57	42	115	298	4.11	97.77
.040 - .049	30	6	11	63	110	1.52	99.29
.050 - .059	9	8		25	42	.58	99.87
.060 - .069	1	1		7	9	.13	100.00
Total	1387	1937	1882	2044	7250	82.76	
Downtime	773	247	326	164	1510	17.24	
Total Time at Site	2160	2184	2208	2208	8760		

TABLE 15

CARBON MONOXIDE DATA  
SITE NUMBER 1  
1983

Month	Hours	Arith. Mean ppm	1 Hr. Max. ppm	No. of Times 1 Hr. Standard (35 ppm) exceeded	8 Hr. Max. ppm	No. of Times 8 Hr. Standard (9 ppm) exceeded
J	516	.91	8.9	-0-	6.5	-0-
F	535	.24	3.9	-0-	2.4	-0-
M	620	.22	4.3	-0-	1.5	-0-
A	685	.28	2.8	-0-	1.5	-0-
M	642	.39	3.1	-0-	2.2	-0-
J	531	.41	3.5	-0-	1.9	-0-
J	426	.24	2.1	-0-	.7	-0-
A	507	.77	2.8	-0-	1.9	-0-
S	444	.44	3.1	-0-	1.5	-0-
O	485	.53	2.3	-0-	1.3	-0-
N	372	1.47	8.7	-0-	3.2	-0-
D	421	1.02	8.1	-0-	4.5	-0-
Annual	6184	.58	8.9	-0-	6.5	-0-

TABLE 16

AMBIENT AIR, CARBON MONOXIDE, PPM  
SITE NUMBER 1  
1983

	Quarter 1	Quarter 2	Quarter 3	Quarter 4	Annual		
1 Hr. Maximum	8.9	3.5	3.1	8.7	8.9		
8 Hr. Maximum	6.5	2.2	1.9	4.5	6.5		
24 Hr. Maximum	2.7	1.4	1.8	2.3	2.7		
Mean.	.91	.41	.77	1.47	.58		
Concentration Range						%	Cum. %
< 2.0	1618	1827	1361	1165	5971	96.56	96.56
2.0 - 2.4	10	23	14	62	109	1.76	98.32
2.5 - 2.9	10	3	1	24	38	.61	98.93
3.0 - 3.4	4	3	1	6	14	.23	99.16
3.5 - 3.9	4	1		5	10	.16	99.32
4.0 - 4.4	6	1		3	10	.16	99.48
4.5 - 4.9	1			4	5	.08	99.56
5.0 - 5.4	1			2	3	.05	99.61
5.5 - 5.9	1			2	3	.05	99.66
6.0 - 6.4	3			2	5	.08	99.74
6.5 - 6.9	2			1	3	.05	99.79
7.0 - 7.4	6				6	.10	99.89
7.5 - 7.9	1				1	.02	99.91
8.0 - 8.4	1			1	2	.03	99.94
8.5 - 8.9	3			1	4	.06	100.00
Total	1671	1858	1377	1278	6184	70.59	
Downtime	489	326	831	930	2576	29.41	
Total Time at Site	2160	2184	2208	2208	8760		



TABLE 17

OZONE DATA  
SITE NUMBER 16  
1983

Month	Hours	Arith. Mean ppm	1 Hour Maximum ppm	No. of Hours 1 Hr. Florida Std. or Federal STD. (0.12 PPM) exceeded
J	695	.017	.058	-0-
F	657	.026	.056	-0-
M	735	.031	.072	-0-
A	714	.030	.087	-0-
M	726	.025	.071	-0-
J	701	.016	.060	-0-
J	0	-	-	-
A	122	.025	.056	-0-
S	610	.015	.052	-0-
O	735	.014	.065	-0-
N	584	.018	.066	-0-
D	434	.011	.047	-0-
Annual	6713	.021	.087	-0-

TABLE 18

AMBIENT AIR, OZONE, PPM  
SITE NUMBER 16  
1983

	Quarter 1	Quarter 2	Quarter 3	Quarter 4	Annual		
1 Hr. Maximum	.072	.087	.056	.066	.087		
8 Hr. Maximum	.062	.071	.049	.053	.071		
24 Hr. Maximum	.041	.044	.040	.030	.044		
Mean.	.031	.030	.025	.018	.021		
Concentration Range			Hours			%	Cum. %
< .010	317	475	266	797	1855	27.63	27.63
.010 - .019	508	483	177	409	1577	23.49	51.12
.020 - .029	493	435	147	300	1375	20.48	71.60
.030 - .039	473	334	102	169	1078	16.06	87.66
.040 - .049	187	248	36	54	525	7.82	95.48
.050 - .059	92	125	4	20	241	3.59	99.07
.060 - .069	16	29		4	49	.73	99.80
.070 - .079	1	8			9	.14	99.94
.080 - .089		4			4	.06	100.00
Total	2087	2141	732	1753	6713	76.63	
Downtime	73	43	1476	455	2047	23.37	
Total Time at Site	2160	2184	2208	2208	8760		

TABLE 19

OZONE DATA  
SITE NUMBER 21  
1983

Month	Hours	Arith. Mean ppm	1 Hour Maximum ppm	No. of Hours 1 Hr. Florida STD. or Federal STD. (0.12 ppm) exceeded
J	665	.020	.054	-0-
F	595	.032	.063	-0-
M	722	.031	.071	-0-
A	711	.034	.088	-0-
M	725	.030	.078	-0-
J	706	.018	.078	-0-
J	131	.015	.060	-0-
A	730	.018	.092	-0-
S	621	.017	.062	-0-
O	708	.017	.064	-0-
N	645	.018	.058	-0-
D	671	.015	.049	-0-
Annual	7630	.021	.092	-0-

TABLE 20

AMBIENT AIR, OZONE, PPM  
SITE NUMBER 21  
1983

	Quarter 1	Quarter 2	Quarter 3	Quarter 4	Annual		
1 Hr. Maximum	.071	.088	.092	.064	.092		
8 Hr. Maximum	.060	.073	.064	.053	.073		
24 Hr. Maximum	.048	.056	.047	.039	.056		
Mean.	.032	.034	.018	.018	.021		
Concentration Range						%	Cum. %
< .010	285	467	579	696	2027	26.57	26.57
.010 - .019	313	322	313	493	1441	18.89	45.46
.020 - .029	429	391	242	509	1571	20.59	66.05
.030 - .039	470	404	202	248	1324	17.35	83.40
.040 - .049	364	278	106	59	807	10.57	93.97
.050 - .059	97	178	27	14	316	4.14	98.11
.060 - .069	23	76	7	5	111	1.46	99.57
.070 - .079	1	21	4		26	.34	99.91
.080 - .089		5	1		6	.08	99.99
.090 - .099			1		1	.01	100.00
Total	1982	2142	1482	2024	7630	87.10	
Downtime	178	42	726	184	1130	12.90	
Total Time at Site	2160	2184	2208	2208	8760		



TABLE 21

OZONE DESIGN VALUES 1980-1983  
SITE NUMBER 16  
WIEBULL DISTRIBUTIONS

Year	Number of Excursions 0.12 ppm	Number of Days Sampled	% Complete	Maximum 1 Hr Avg. ppm	2nd Max. 1 Hr Avg. ppm	Annual Design Value ppm	Three Year Design Value ppm
1980	-0-	292	79.78	.098	.083	.097	N.A.
1981	-0-	318	87.12	.095	.089	.095	N.A.
1982	-0-	352	96.44	.080	.078	.084	.095
1983	-0-	298	81.64	.087	.085	.088	.088

TABLE 22

OZONE DESIGN VALUES 1973-1983  
 SITE NUMBER 21  
 WIEBULL DISTRIBUTIONS

Year	Number of Excursions 0.12 ppm	Number of Days Sampled	% Complete	Maximum 1 Hr Avg. ppm	2nd Max. 1 Hr Avg. ppm	Annual Design Value ppm	Three Year Design Value ppm
1973	-0-	116	31.78	.111	.104	.130	N.A.
1974	-0-	363	99.45	.077	.074	.078	.106
1975	-0-	331	90.68	.104	.086	.097	.101
1976	3	270	73.97	.148	.142	.154	.123
1977	-0-	323	88.49	.106	.101	.106	.122
1978	-0-	138	37.81	.075	.069	.079	.128
1979	-0-	256	70.14	.082	.076	.081	.092
1980	-0-	299	81.69	.110	.102	.109	.101
1981	-0-	348	95.34	.103	.089	.098	.102
1982	-0-	354	96.99	.122	.080	.106	.106
1983	-0-	332	90.96	.092	.088	.091	.101

TABLE 23

SULFUR DIOXIDE DATA  
 SITE NUMBER 22  
 1983

Month	Hours	Arith. Mean ppm	1 Hr Max. ppm	3 Hr Max. ppm	No. of Times 3 Hr Std. Exceeded (0.5 ppm)	24 Hr Max. ppm	No. of Times 24 Hr Std. Exceeded (0.1 ppm)
J	543	.003	.018	.018	-0-	.015	-0-
F	630	.0004	.011	.006	-0-	.003	-0-
M	732	.0004	.015	.013	-0-	.005	-0-
A	709	.0002	.007	.007	-0-	.003	-0-
M	740	.0003	.021	.017	-0-	.003	-0-
J	679	.0001	.008	.007	-0-	.002	-0-
J	740	.0005	.035	.025	-0-	.005	-0-
A	728	.0003	.018	.009	-0-	.001	-0-
S	680	.0002	.010	.008	-0-	.002	-0-
O	731	.00004	.007	.005	-0-	.001	-0-
N	711	.00002	.003	.002	-0-	.0003	-0-
D	737	.00003	.013	.007	-0-	.0009	-0-
Annual	8360	.0005	.035	.025	-0-	.015	-0-

TABLE 24

AMBIENT AIR, SULFUR DIOXIDE, PPM  
 SITE NUMBER 22  
 1983

	Quarter 1	Quarter 2	Quarter 3	Quarter 4	Annual		
1 Hr Maximum	.018	.021	.035	.013	.035		
8 Hr Maximum	.017	.010	.014	.003	.017		
24 Hr Maximum	.015	.003	.005	.001	.015		
Arithmetic Mean	.003	.0003	.0005	.00004	.0005		
Concentration Range			Hours			%	Cum. %
< .010	1853	2125	2130	2178	8286	99.12	99.12
.010 - .014	37	1	8	1	47	.57	99.69
.015 - .019	15	1	5		21	.25	99.94
.020 - .024		1	1		2	.02	99.96
.025 - .029							
.030 - .034			2		2	.02	99.98
.035 - .039			2		2	.02	100.00
Total	1905	2128	2148	2179	8360	95.43	
Downtime	255	56	60	29	400	4.57	
Total Time at Site	2160	2184	2208	2208	8760		



#### IV

#### AIR QUALITY INDEX

The Environmental Protection Agency (EPA) has developed a uniform standardized daily air quality reporting index, called the Pollutant Standard Index (PSI), locally called the Air Quality Index (AQI), to be used by State and local agencies. The use of this index for reporting air quality was made mandatory on May 10, 1979 in the Federal Register/Vol. 44, No 92/Part 58, Subpart E, Appendix G.

This index is dependent upon measured concentrations of the five pollutants which have been assigned National Ambient Air Quality Standards (NAAQS), Federal Episode Criteria, and Significant Harm Levels; i.e., total suspended particulate, carbon monoxide, sulfur dioxide, nitrogen dioxide, and ozone. The index converts air pollution concentrations to a normalized number on a scale of zero to five hundred, with the National Ambient Air Quality Standard for each pollutant being assigned the value of 100. This approach is believed to be easier for the public to understand than a report of actual pollutant concentrations. Index values are calculated for each of the five pollutants. The highest of these is the report value. Five descriptor words have been chosen to depict daily air quality: "good" (0-50), "moderate" (51-100), "unhealthful" (101-200), "very unhealthful" (201-300), and "hazardous" (301-500). If pollutant concentration warrant, the AQI report is expanded to include identification of the problem pollutant, cautionary statements and generalized health effects.

Adoption of the AQI by many pollution control organizations has reduced the confusion previously encountered due to the existances of many different indices. AQI has several advantages: (1) it is simple and can be easily understood by the public, (2) it can accommodate new pollutants, (3) it is based on a reasonable scientific premise, (4) it relates to National Ambient Air Quality Standards, Federal Episode Criteria, and Significant Harm Levels, and (5) it exhibits day to day variations.

The Palm Beach County Health Department samples for the above pollutants and has utilized the Air Quality Index since April 28, 1976. A typical air quality report for Palm Beach County would be "The Air Quality Index is 31. The ambient air quality is within the good range." Index advisories are issued to local newspapers, televisions and radio stations each morning and afternoon Monday through Friday. Table 25 gives a statistical analysis of the monthly morning and afternoon values. As noted, TSP or Ozone concentrations accounted for all values in the moderate range.

TABLE 25

AIR QUALITY INDEX  
WEST PALM BEACH  
1983

## A.M. DATA

Month	Days		A Q I			
	Good	Moderate	Maximum	Minimum	Mean.	SD.
Jan.	21	-0-	39	10	22	7
Feb.	20	-0-	34	12	24	6
March	21	2	59	14	29	10
April	20	1	59	15	27	11
May	21	-0-	34	14	24	5
June	22	-0-	31	11	19	6
July	17	3	69	12	32	15
Aug.	23	-0-	44	15	23	7
Sept.	20	1	53	12	27	11
Oct.	21	-0-	26	8	14	5
Nov.	19	-0-	35	14	23	6
Dec.	21	-0-	27	11	19	5
Year	246	7	69	8	24	9

## P.M. DATA

Month	Days		A Q I			
	Good	Moderate	Maximum	Minimum	Mean.	SD.
Jan.	21	-0-	40	15	28	7
Feb.	20	-0-	50	25	34	6
March	22	1	58	14	33	9
April	18	3	73	17	41	13
May	20	1	78	19	36	13
June	22	-0-	47	10	27	8
July	17	4	69	11	35	15
Aug.	22	1	79	18	34	13
Sept.	20	1	53	18	29	8
Oct.	20	1	54	12	24	12
Nov.	18	1	51	19	29	7
Dec.	21	-0-	42	10	26	7
Year	241	13	79	10	31	11

Of the values in the Moderate category, T.S.P. values were responsible for (7) morning and (2) afternoon A.Q.I.'s. All others in the moderate range were related to ozone concentrations. No A.Q.I. values above the moderate zone were reported.



## EMISSION INVENTORY

## INTRODUCTION

Air pollution evolved as an undesired by-product of the technological advancement of our modern society. On the other hand, this modern society also developed technological methods to prevent and control these atmospheric emissions of air pollutants.

Defining air pollution problem areas by determining the actual sources and components of air pollution is a prime step towards abating air pollution, and subsequently attaining and maintaining national ambient air quality standards (NAAQS). The emission inventory accomplishes this crucial step by addressing the sources, types and quantities of air pollution emissions.

A comprehensive emission inventory is an essential tool for any Air Pollution regulatory agency. The inventory provides information for the design of an air sampling and analysis program, identifies the relative contribution of the various pollution sources, offers data for the development of control strategies, and makes this information available for our regional planning authorities.

A number of factors used in compiling the emission inventory, such as reported fuel usage figures, fuel composition, process information and especially emission factors, are somewhat limited in preciseness due to the state-of-the-art knowledge available. Never-the-less, the emission inventory, by nature of its magni-



tude, still yields results with adequate accuracy for the purpose and intent of our county air pollution program.

The emission inventory presented in this report represents calculated emissions from major point sources and generalized acceptable estimates of emissions from area sources (Tables 26 and 27, respectively).

#### MAJOR POINT SOURCES

Palm Beach County has 23 major air pollution sources which emit a significant emission tonage of one or more types of air pollutants. Table 26 lists these sources by name and details air emissions by pollutant type in tons per year for 1983. A constantly unpredictable economy has caused a reduction in emission tonage for certain sources due to reduced activity, whereas other sources have actually increased their activity, hence increased their emission tonage above prior years. Data used to compile and quantify the emission levels for each source was gathered using specific data procurement techniques, such as annual operating reports, process material balances, engineering appraisals, and special questionnaires.

#### AREA SOURCES

Palm Beach County has many sources which emit small quantities of air pollutants, and collectively their impact is significant to our atmosphere. Therefore, these sources must be accounted for in the emission inventory. Table 27 lists major area source categories and details respective air emissions by types in tons per year for 1983. The object of area source calculations is to obtain an accurate estimate of this collective

contribution on total emissions. Such an estimate may never be exact due to the difficulties of determining the emissions from every source too small and/or too numerous to be surveyed individually. Ultimately, however, this area source estimate becomes more valid as it is applied to a large number of sources since prescribed techniques were used in conjunction with the activity levels associated with each source category.

References for emission factors utilized are:

- (1) Compilation of Air Pollution Emission Factors, AP-42, Supplements 1-13, Third Edition, August 1982, published by the U. S. Environmental Protection Agency, Office of Air and Waste Management, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, 27711.
- (2) Guidelines for Air Quality Maintenance Planning and Analysis, Volume 7: Projecting County Emissions, EPA-450/4-74-008, Second Edition, January 1975, U. S. Environmental Protection Agency, Office of Air and Waste Management, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, 27711.
- (3) National Emissions Data System (NEDS) Source Classification Code and Emission Factor Listing, January 1979, U. S. Environmental Protection Agency, Office of Air Quality Planning and Standards, Monitoring and Data Analysis Division,

National Air Data Branch, Research Triangle Park, North Carolina, 27711.

- (4) Procedures for the Preparation of Emission Inventories for Volatile Organic Compounds, Volume I, EPA-450/2-77-028, December 1977, U. S. Environmental Protection Agency, Office of Air and Water Management, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, 27711.
- (5) Mobile Source Emission Factors (For Low-Altitude Areas Only), EPA-400/9-78-006, March 1978, U. S. Environmental Protection Agency, Office of Air and Waste Management, Office of Transportation and Land Use Policy, Washington, D. C. 20460.
- (6) Revisions to the State Implementation Plan (SIP) for Palm Beach County, Florida, April 1983, Metropolitan Planning Organization of Palm Beach County.
- (7) Procedures for Emission Inventory Preparation, Volumes I-IV, EPA-450/4-81-026A, September 1981, United States Environmental Protection Agency, Office of Air Quality Planning and Standards, Research Triangle Park, North Carolina, 27711.

TABLE 26

1983 SUMMARY  
 AIR POLLUTION EMISSIONS  
 PALM BEACH COUNTY, FLORIDA  
 TONS/YEAR  
 POINT SOURCES

A.P.I.S. No 50-50	Source Name	Parti- culates	SO <sub>2</sub>	SO <sub>3</sub>	CO	CH <sub>4</sub>	NO <sub>2</sub>	VOC	TOTAL
0005	Gulf & Western Food Co.	851.3	798.1	.3	667.9	253.5	418.2	413.2	3402.5
0011	Florida Asphalt Co.	.5	11.5	.2	.8		3.2		16.2
0015	Boca Raton Hotel & Club	1.8	55.9	.7	1.0	.1	10.2	4.4	74.1
0016	Atlantic Sugar Assoc.	382.5	400.3	.9	334.9	127.0	219.0	206.7	1671.3
0018	Quaker Oats Co.	1.2	25.2	.4	3.0		11.8	.2	41.9
0019	Osceola Farms Co.	432.5	462.9	1.4	353.2	133.7	227.7	218.2	1829.6
0021	Pratt & Whitney	69.8	78.1		1196.2	19.8	175.5	326.6	1866
0026	Sugar Cane Growers Corp.	545.6	1615.9	.3	593.1	225.3	363.7	367.3	3711.2
0042	Florida Power & Light	383.3	3904.2	71.9	610.1	10.6	8341.1	35.9	13357.1
0045	Lake Worth Utilities	19.6	26.7	.3	148.2	21.6	698.1	27.4	941.9
0046	Eagle Cement	1.1							1.1
0050	Florida Sugar Refinery	6.9	96.9	1.2	1.3	.2	13.8	.1	120.4
0061	U. S. Sugar Corp-Bryant	409.6	546.7	.4	514.8	195.5	314.5	318.9	2300.4
0073	Talisman Sugar Corp.	203.4	298.5	.2	283.5	107.7	172.5	175.6	1241.4
0081	A. G. Holley Hospital	1.4	6.6	.1	1.4	.2	1.7	.3	11.7
0084	Eastern Cement	14.6							14.6
0087	Ranger Construction Ind.	15.1	29.8	.4	1.8		7.4	.1	54.6
0088	City of Pahokee	9.8	3.7		14.0	1.6	4.3	2.6	36.0
0155	F. H. Foster Oil Corp.							14.6	14.6
0158	Cooper Oil Co.							21.9	21.9
0161	Howell Oil							21.3	21.3
0162	Charles Brown Oil							31.3	31.3
0163	Berner Oil Co.							18.4	18.4
Point Source Totals		3350.0	8361.0	78.7	4725.2	1096.9	10982.7	2205.0	30799.5



TABLE 27

1983 SUMMARY  
 AIR POLLUTION EMISSIONS  
 PALM BEACH COUNTY, FLORIDA  
 TON/YEAR  
 AREA SOURCES

	Particulates	SO <sub>2</sub>	SO <sub>3</sub>	CO	CH <sub>4</sub>	NO <sub>2</sub>	VOC	Aldehydes	Organic Acids	Totals
Mobile Sources:										
Highway/Off Highway	1543	947		147900	1422	10562	13568	45	43	176120
Aircraft	13.1	51.6		5205.9	15.8	404.5	371.2			6062.1
Vessels	17.1	183		3.7	.4	32.7	3.3			240.2
Railroads	35.4	80.7		184.1	2.8	524.1	130.3	7.8	9.9	975.1
Fuel Combustion:										
Miscellaneous	34.5	622.7	8.8	93.4	17.6	395.2	17.6			1189.8
Mineral Products										
Concrete Batching	55.6									55.6
Solid Waste Disposal:										
Incineration	11.0	3.2		13.9	3.1	11.1	3.4			45.7
Open Burning	223.2			1838.1	119.7		195.4			2376.4
Sugar Cane Fields:										
Burning	7886			76047	5708		9313			98954
Volatile Organic Emissions:										
Storage & Marketing of										
Petrol Prod.							1690.5			1690.5
Industrial Processes							146			146
Industrial Surface Ctg.							212.1			212.1
Non-Indus. Surface Ctg.							854			854
Other Solvent Use							3695.2			3695.2
Area Source Totals	9818.9	1888.2	8.8	231286.1	7289.4	12019.6	30200.0	52.8	52.9	292616.7
Point Source Totals	3350.0	8361.0	78.7	4725.2	1096.9	10982.7	2205.0	0	0	30799.5
Grand Totals	13168.9	10249.2	87.5	236011.3	8386.3	23002.3	32405.0	52.8	52.9	323416.2



